
ON THE VERTEX FOLKMAN NUMBERS

$$F_V(\underbrace{2, \dots, 2}_R; R-1) \text{ AND } F_V(\underbrace{2, \dots, 2}_R; R-2)$$

NEDYALKO NENOV

For a graph G the symbol $G \xrightarrow{a} (a_1, \dots, a_r)$ means that in every r -coloring of the vertices of G , for some $i \in \{1, 2, \dots, r\}$ there exists a monochromatic a_i -clique of color i . The vertex Folkman numbers

$$F_v(a_1, \dots, a_r; q) = \min\{|V(G)| : G \xrightarrow{a} (a_1, \dots, a_r) \text{ and } K_q \not\subseteq G\}$$

are considered. We prove that

$$F_v(\underbrace{2, \dots, 2}_r; r-1) = r+7, \quad r \geq 6 \quad \text{and} \quad F_v(\underbrace{2, \dots, 2}_r; r-2) = r+9, \quad r \geq 8.$$

Keywords: Folkman graphs, Folkman numbers

2000 Math. Subject Classification: 05C55

1. INTRODUCTION

We consider only finite, non-oriented graphs without loops and multiple edges. We call a p -clique of the graph G a set of p vertices, each two of which are adjacent. The largest positive integer p such that the graph G contains a p -clique is denoted by $\text{cl}(G)$. In this paper we shall also use the following notation:

- $V(G)$ is the vertex set of the graph G ;
- $E(G)$ is the edge set of the graph G ;

- \overline{G} is the complement of G ;
- $G[V]$, $V \subseteq V(G)$ is the subgraph of G induced by V ;
- $G - V$, $V \subseteq V(G)$ is the subgraph of G induced by $V(G) \setminus V$;
- $\alpha(G)$ is the vertex independence number of G ;
- $\chi(G)$ is the chromatic number of G ;
- $f(G) = \chi(G) - \text{cl}(G)$;
- K_n is the complete graph on n vertices;
- C_n is the simple cycle on n vertices.

Let G_1 and G_2 be two graphs without common vertices. We denote by $G_1 + G_2$ the graph G for which $V(G) = V(G_1) \cup V(G_2)$ and $E(G) = E(G_1) \cup E(G_2) \cup E'$, where $E' = \{[x, y] : x \in V(G_1), y \in V(G_2)\}$.

The Ramsey number $R(p, q)$ is the smallest natural n such that for every n -vertex graph G either $\text{cl}(G) \geq p$ or $\alpha(G) \geq q$. An exposition of the results on the Ramsey numbers is given in [25]. In Table 1.1 we list the known Ramsey numbers $R(p, 3)$ (see [25]).

p	3	4	5	6	7	8	9	10
$R(p, 3)$	6	9	14	18	23	28	36	40–43

Table 1.1: The known Ramsey numbers

Definition. Let a_1, \dots, a_r be positive integers. We say that the r -coloring

$$V(G) = V_1 \cup \dots \cup V_r, \quad V_i \cap V_j = \emptyset, \quad i \neq j$$

of the vertices of the graph G is (a_1, \dots, a_r) -free, if V_i does not contain an a_i -clique for each $i \in \{1, \dots, r\}$. The symbol $G \xrightarrow{v} (a_1, \dots, a_r)$ means that there is no (a_1, \dots, a_r) -free coloring of the vertices of G .

Let a_1, \dots, a_r and q be natural numbers. Define

$$H_v(a_1, \dots, a_r; q) = \{G : G \xrightarrow{v} (a_1, \dots, a_r) \text{ and } \text{cl}(G) < q\},$$

$$F_v(a_1, \dots, a_r; q) = \min\{|V(G)| : G \in H_v(a_1, \dots, a_r; q)\}.$$

The graph $G \in H_v(a_1, \dots, a_r; q)$ is said to be an extremal graph in $H_v(a_1, \dots, a_r; q)$, if $|V(G)| = F_v(a_1, \dots, a_r; q)$.

It is clear that $G \xrightarrow{v} (a_1, \dots, a_r)$ implies $\text{cl}(G) \geq \max\{a_1, \dots, a_r\}$. Folkman [3] proved that there exists a graph G such that $G \xrightarrow{v} (a_1, \dots, a_r)$ and $\text{cl}(G) = \max\{a_1, \dots, a_r\}$. Therefore

$$F_v(a_1, \dots, a_r; q) \text{ exists} \iff q > \max\{a_1, \dots, a_r\}. \quad (1.1)$$

The numbers $F_v(a_1, \dots, a_r; q)$ are called vertex Folkman numbers.

If a_1, \dots, a_r are positive integers, $r \geq 2$ and $a_i = 1$ then it is easily seen that

$$G \xrightarrow{v} (a_1, \dots, a_i, \dots, a_r) \iff G \xrightarrow{v} (a_1, \dots, a_{i-1}, a_{i+1}, a_r).$$

Thus it suffices to consider only such numbers $F_v(a_1, \dots, a_r; q)$ for which $a_i \geq 2$, $i = 1, \dots, r$. In this paper we consider the vertex Folkman numbers $F_v(2, \dots, 2; q)$. Set

$$\underbrace{(2, \dots, 2)}_r = (2_r) \quad \text{and} \quad F_v(\underbrace{2, \dots, 2}_r; q) = F_v(2_r; q).$$

By (1.1),

$$F_v(2_r; q) \text{ exists} \iff q \geq 3. \quad (1.2)$$

It is clear that

$$G \xrightarrow{v} (2_r) \iff \chi(G) \geq r + 1. \quad (1.3)$$

Since $K_{r+1} \xrightarrow{v} (2_r)$ and $K_r \not\xrightarrow{v} (2_r)$, we have

$$F_v(2_r; q) = r + 1 \quad \text{if} \quad q \geq r + 2.$$

In [2] Dirac proved the following result.

Theorem 1.1. ([2]) *Let G be a graph such that $\chi(G) \geq r + 1$ and $\text{cl}(G) \leq r$. Then*

- (a) $|V(G)| \geq r + 3$;
- (b) *If $|V(G)| = r + 3$, then $G = K_{r-3} + C_5$.*

According to (1.3), Theorem 1.1 admits the following equivalent form:

Theorem 1.2. *Let $r \geq 2$ be a positive integer. Then*

- (a) $F_v(2_r; r + 1) = r + 3$;
- (b) $K_{r-2} + C_5$ *is the only extremal graph in $H_v(2_r; r + 1)$.*

In [14] Łuczak, Ruciński and Urbański defined for arbitrary positive integers a_1, \dots, a_r the numbers

$$m = \sum_{i=1}^r (a_i - 1) + 1 \quad \text{and} \quad p = \max\{a_1, \dots, a_r\}. \quad (1.4)$$

They proved the following extension of Theorem 1.2.

Theorem 1.3. ([14]) *Let a_1, \dots, a_r be positive integers and m and p be defined by (1.4). Let $m \geq p + 1$. Then*

- (a) $F_v(a_1, \dots, a_r; m) = m + p;$
- (b) $K_{m-p-1} + \overline{C}_{2p+1}$ is the only extremal graph in $H_v(a_1, \dots, a_r; m).$

For another extension of Theorem 1.1 see [21].

From (1.1) it follows that the numbers $F_v(a_1, \dots, a_r; m - 1)$ exist if and only if $m \geq p + 2$. The exact values of all numbers $F_v(a_1, \dots, a_r; m - 1)$ for which $p = \max\{a_1, \dots, a_r\} \leq 4$ are known. A detailed exposition of these results was given in [13] and [23]. We do not know any exact values of $F_v(a_1, \dots, a_r; m - 1)$ in the case when $\max\{a_1, \dots, a_r\} \geq 5$. Here we shall note only the values $F_v(a_1, \dots, a_r; m - 1)$ when $a_1 = a_2 = \dots = a_r = 2$, i.e. of the numbers $F_v(2_r; r)$. From (1.2) these numbers exist if and only if $r \geq 3$. If $r = 3$ and $r = 4$ we have that

$$F_v(2_3; 3) = 11; \tag{1.5}$$

$$F_v(2_4; 4) = 11. \tag{1.6}$$

The inequality $F_v(2_3; 3) \leq 11$ was proved in [15] and the opposite inequality $F_v(2_3; 3) \geq 11$ was proved in [1]. The equality (1.6) was proved in [18] (see also [19]). If $r \geq 5$ we have the following result.

Theorem 1.4. ([17], see also [24]) *Let $r \geq 5$. Then:*

- (a) $F_v(2_r; r) = r + 5;$
- (b) $K_{r-5} + C_5 + C_5$ is the only extremal graph in $H_v(2_r; r).$

Theorem 1.4(a) was proved also in [8] and [14].

According to (1.2), the number $F_v(2_r; r - 1)$ exists if and only if $r \geq 4$. In [17] we proved that

$$F_v(2_r; r - 1) = r + 7 \quad \text{if } r \geq 8. \tag{1.7}$$

In this paper we improve (1.7) by proving the following result:

Theorem 1.5. *Let $r \geq 4$ be an integer. Then:*

- (a) $F_v(2_r; r - 1) \geq r + 7;$
- (b) $F_v(2_r; r - 1) = r + 7, \quad \text{if } r \geq 6;$
- (c) $F_v(2_5; 4) \leq 16.$

In [9] Jensen and Royle showed that

$$F_v(2_4; 3) = 22. \tag{1.8}$$

We see from Theorem 1.5 and (1.8) that $F_v(2_5; 4)$ is the only unknown number of the kind $F(2_r; r - 1)$ ¹.

From (1.2) it follows that the Folkman number $F(2_r; r - 2)$ exists if and only if $r \geq 5$. In [16] we proved that $F_v(2_r; r - 2) = r + 9$ if $r \geq 11$. In this paper we improve this result as follows:

Theorem 1.6. *Let $r \geq 5$ be an integer. Then:*

- (a) $F_v(2_r; r - 2) \geq r + 9$;
- (b) $F_v(2_r; r - 2) = r + 9$, if $r \geq 8$.

The numbers $F_v(2_r; r - 2)$, $5 \leq r \leq 7$, are unknown.

2. AUXILIARY RESULTS

Let G be an arbitrary graph. Define

$$f(G) = \chi(G) - \text{cl}(G).$$

Lemma 2.1. *Let G be a graph such that $f(G) \leq 2$. Then*

$$|V(G)| \geq \chi(G) + 2f(G).$$

Proof. Since $\chi(G) \geq \text{cl}(G)$, we have $f(G) \geq 0$. For $f(G) = 0$ the inequality is trivial. Let $f(G) = 1$ and $\chi(G) = r + 1$. Then $\text{cl}(G) = r$. Note that $r \geq 2$ because of $\chi(G) \neq \text{cl}(G)$. By (1.3) we have $G \in H_v(2_r; r + 1)$. Thus, from Theorem 1.2(a) it follows that $|V(G)| \geq r + 3 = 2f(G) + \chi(G)$. Let $f(G) = 2$ and $\chi(G) = r + 1$. Then $\text{cl}(G) = r - 1$. Since $\chi(G) \neq \text{cl}(G)$, $\text{cl}(G) = r - 1 \geq 2$, i.e. $r \geq 3$. From Theorem 1.4(a), (1.5) and (1.6) we obtain that $|V(G)| \geq r + 5 = \chi(G) + 2f(G)$. This completes the proof of Lemma 2.1. \square

Let $G = G_1 + G_2$. Obviously,

$$\chi(G) = \chi(G_1) + \chi(G_2); \tag{2.1}$$

$$\text{cl}(G) = \text{cl}(G_1) + \text{cl}(G_2). \tag{2.2}$$

Hence,

$$f(G) = f(G_1) + f(G_2). \tag{2.3}$$

¹Meanwhile, it has been proved that $F_v(2_5; 4) = 16$, see J. Lathrop, S. Radziszowski, Computing the Folkman Number $F_v(2, 2, 2, 2, 2; 4)$, Journal of Combinatorial Mathematics and Combinatorial Computing, 78 (2011), 213–222.

A graph G is said to be vertex-critical chromatic if $\chi(G - v) < \chi(G)$ for all $v \in V(G)$. We shall use the following result in the proof of Theorem 1.6.

Theorem 2.1. ([4], see also [5]) *Let G be a vertex-critical chromatic graph and $\chi(G) \geq 2$. If $|V(G)| < 2\chi(G) - 1$, then $G = G_1 + G_2$, where $V(G_i) \neq \emptyset$, $i = 1, 2$.*

Remark. In the original statement of Theorem 2.1 the graph G is supposed to be edge-critical chromatic (and not vertex-critical chromatic). Since each vertex-critical chromatic graph G contains an edge-critical chromatic subgraph H such that $\chi(G) = \chi(H)$ and $V(G) = V(H)$, the above statement is equivalent to the original one. It is also more convenient for the proof of Theorem 1.6.

Let G be a graph and $A \subseteq V(G)$ be an independent set of vertices of the graph G . It is easy to see that

$$G \xrightarrow{v} (2_r), r \geq 2 \Rightarrow G - A \xrightarrow{v} (2_{r-1}). \quad (2.4)$$

Lemma 2.2. *Let $G \in H_v(2_r; q)$, $q \geq 3$ and $|V(G)| = F_v(2_r; q)$. Then*

- (a) G is a vertex-critical $(r + 1)$ -chromatic graph;
- (b) If $q < r + 3$, then $\text{cl}(G) = q - 1$.

Proof. By (1.3), $\chi(G) \geq r + 1$. Assume that (a) is false. Then there would exist $v \in V(G)$ such that $\chi(G - v) \geq r + 1$. According to (1.3), $G - v \in H_v(2_r; q)$. This contradicts the equality $|V(G)| = F_v(2_r; q)$.

Assume that (b) is false, i.e. $\text{cl}(G) \leq q - 2$. Then from $q < r + 3$ it follows that $\text{cl}(G) < r + 1$. Since $\chi(G) \geq r + 1$ there are $a, b \in V(G)$ such that $[a, b] \notin E(G)$. Consider the subgraph $G_1 = G - \{a, b\}$. We have $r \geq 2$, because $\chi(G) \neq \text{cl}(G)$. Thus, from (2.4) and $\text{cl}(G) \leq q - 2$ it follows that $G_1 \in H_v(2_{r-1}; q - 1)$. Obviously, $G_1 \in H_v(2_{r-1}; q - 1)$ leads to $K_1 + G_1 \in H_v(2_r; q)$. This contradicts the equality $|V(G)| = F_v(2_r; q)$, because $|V(K_1 + G_1)| = |V(G)| - 1$. Lemma 2.2 is proved. \square

Lemma 2.3. *Let $G \in H_v(2_r; q)$, $r \geq 2$. Then*

$$|V(G)| \geq F_v(2_{r-1}; q) + \alpha(G).$$

Proof. Let $A \subseteq V(G)$ be an independent set such that $|A| = \alpha(G)$. Consider the subgraph $G_1 = G - A$. According to (2.4), $G_1 \in H_v(2_{r-1}; q)$. Hence $|V(G_1)| \geq F_v(2_{r-1}; q)$. Since $|V(G)| = |V(G_1)| + \alpha(G)$, Lemma 2.3 is proved. \square

We shall use also the following three results:

$$F_v(2, 2, p; p + 1) \geq 2p + 4, \quad \text{see [20]}; \quad (2.5)$$

$$F_v(2, 2, 4; 5) = 13, \quad \text{see [22]}. \quad (2.6)$$

Theorem 2.2. ([12]) *Let G be a graph, $\text{cl}(G) \leq p$ and $|V(G)| \geq p+2$, $p \geq 2$. Let G also possess the following two properties:*

- (i) $G \not\rightarrow (2, 2, p)$;
- (ii) *If $V(G) = V_1 \cup V_2 \cup V_3$ is a $(2, 2, p)$ -free 3-coloring, then $|V_1| + |V_2| \leq 3$.*

Then $G = K_1 + G_1$.

3. AN UPPER BOUND FOR THE NUMBERS $F_v(2r; q)$

Consider the graph P whose complementary graph \overline{P} is depicted in Figure 1. This graph is a well-known construction of Greenwood and Gleason [6], which




Figure 1: Graph \overline{P}

shows that $R(5, 3) \geq 14$, since $|V(P)| = 13$ and

$$\alpha(P) = 2; \tag{3.1}$$

$$\text{cl}(P) = 4 \quad (\text{see [6]}). \tag{3.2}$$

From $|V(P)| = 13$ and (3.1) it follows that $\chi(P) \geq 7$. Since $\{v_1\} \cup \{v_2, v_3\} \cup \dots \cup \{v_{12}, v_{13}\}$ is a 7-chromatic partition of $V(P)$, we have

$$\chi(P) = 7. \tag{3.3}$$

Let r and s be non-negative integers and $r \geq 3s + 6$. Define

$$\tilde{P} = K_{r-3s-6} + P + \underbrace{C_5 + \dots + C_5}_s.$$

From (2.1), (2.2), (3.2) and (3.3) we obtain that $\chi(\tilde{P}) = r + 1$ and $\text{cl}(\tilde{P}) = r - s - 2$. By (1.3), it follows that $\tilde{P} \in H_v(2_r; r - s - 1)$ and thus

$$F_v(2_r; r - s - 1) \leq |V(\tilde{P})|.$$

Since $|V(\tilde{P})| = r + 2s + 7$, we proved the following

Theorem 3.1. *Let r and s be non-negative integers and $r \geq 3s + 6$. Then*

$$F_v(2_r; r - s - 1) \leq r + 2s + 7.$$

Remark. Since $r \geq 3s + 6$ we have $r - s - 1 > 2$. Thus, according to (1.2), the numbers $F_v(2_r; r - s - 1)$ exist.

4. PROOF OF THEOREM 1.5

Proof of Theorem 1.5(a) Let $G \in H_v(2_r; r - 1)$. We need to show that $|V(G)| \geq r + 7$. From Lemma 2.3 we have

$$|V(G)| \geq F_v(2_{r-1}; r - 1) + \alpha(G).$$

By (1.5), (1.6) and Theorem 1.4(a) we deduce $F_v(2_{r-1}; r - 1) \geq r + 4$. Hence

$$|V(G)| \geq r + 4 + \alpha(G). \quad (4.1)$$

We prove the inequality $|V(G)| \geq r + 7$ by induction with respect to r . From Table 1.1 we see that

$$R(r - 1, 3) < r + 6 \text{ if } r = 4 \text{ or } r = 5. \quad (4.2)$$

Obviously, from $G \in H_v(2_r; r - 1)$ it follows that $\chi(G) \neq \text{cl}(G)$. Thus, $\alpha(G) \geq 2$. From (4.1) we obtain $|V(G)| \geq r + 6$. From this inequality and (4.2) we see that $|V(G)| > R(r - 1, 3)$ if $r = 4$ or $r = 5$. Since $\text{cl}(G) < r - 1$, it follows that $\alpha(G) \geq 3$. Now from (4.1) we obtain that $|V(G)| \geq r + 7$ if $r = 4$ or $r = 5$.

Let $r \geq 6$. We shall consider separately two cases:

Case 1. $G \not\rightarrow (2, 2, r - 2)$. From Theorem 2.2 we see that only following two subcases are possible:

Subcase 1a. $G = K_1 + G_1$. From $G \in H_v(2_r, r - 1)$ it follows that $G_1 \in H_v(2_{r-1}; r - 2)$. By the induction hypothesis, $|V(G_1)| \geq r + 6$. Therefore, $|V(G)| \geq r + 7$.

Subcase 1b. There is a $(2, 2, r - 2)$ -free 3-coloring $V(G) = V_1 \cup V_2 \cup V_3$ such that $|V_1| + |V_2| \geq 4$. Let us consider the subgraph $\tilde{G} = G[V_3]$. By assumption \tilde{G} does not contain an $(r - 2)$ -clique, i.e. $\text{cl}(\tilde{G}) < r - 2$. Since V_1 and V_2 are

independent sets and $G \xrightarrow{v} (2_r)$, it follows from (2.4) that $\tilde{G} \xrightarrow{v} (2_{r-2})$. Thus, $\tilde{G} \in H_v(2_{r-2}; r-2)$. By (1.6) and Theorem 1.4(a), $|V(\tilde{G})| \geq r+3$. As $|V_1|+|V_2| \geq 4$, we have $|V(G)| \geq r+7$.

Case 2. $G \xrightarrow{v} (2, 2, r-2)$. Since $\text{cl}(G) < r-1$, $G \in H_v(2, 2, r-2; r-1)$. From (2.5) it follows that $|V(G)| \geq 2(r-2) + 4 = 2r$. Hence, if $2r \geq r+7$, i.e. $r \geq 7$, then $|V(G)| \geq r+7$. Let $r = 6$. Then $G \in H_v(2, 2, 4; 5)$. By (2.6) we conclude that $|V(G)| \geq 13$.

Proof of Theorem 1.5(b) Let $r \geq 6$. According to Theorem 1.5(a) we have $F_v(2_r; r-1) \geq r+7$. From Theorem 3.1 ($s = 0$) we obtain the opposite inequality $F_v(2_r; r-1) \leq r+7$.

Proof of Theorem 1.5(c) There is a 16-vertex graph G such that $\alpha(G) = 3$ and $\text{cl}(G) = 3$, because $R(4, 4) = 18$ (see [6]). From $|V(G)| = 16$ and $\alpha(G) = 3$ obviously it follows that $\chi(G) \geq 6$. By (1.3), $G \xrightarrow{v} (2_5)$. So, $G \in H_v(2_5; 4)$. Hence $F_v(2_5; 4) \leq |V(G)| = 16$.

Theorem 1.5 is proved. □

Corollary 4.1 *Let G be a graph such that $f(G) \leq 3$. Then*

$$|V(G)| \geq \chi(G) + 2f(G).$$

Proof. If $f(G) \leq 2$, then Corollary 4.1 follows from Lemma 2.1. Let $f(G) = 3$ and $\chi(G) = r+1$, then $\text{cl}(G) = r-2$. Since $\chi(G) \neq \text{cl}(G)$, it follows that $\text{cl}(G) \geq 2$. Thus, $r \geq 4$. By (1.3) we get $G \in H_v(2_r; r-1)$. From Theorem 1.5(a) we obtain $|V(G)| \geq r+7 = \chi(G) + 2f(G)$. □

Remark. In $H_v(2_r; r-1)$, $r \geq 8$, there are more than one extremal graph. For instance, in $H_v(2_8; 7)$ besides $K_2 + P$ (see Theorem 3.1), the graph $C_5 + C_5 + C_5$ is extremal, too.

5. PROOF OF THEOREM 1.6

Proof of Theorem 1.6(a) Let $G \in H_v(2_r; r-2)$. We need to show that $|V(G)| \geq r+9$. From Lemma 2.3 we have

$$|V(G)| \geq F_v(2_{r-1}; r-2) + \alpha(G).$$

By Theorem 1.5(a), $F_v(2_{r-1}; r-2) \geq r+6$. Thus,

$$|V(G)| \geq r+6 + \alpha(G). \tag{5.1}$$

We prove the inequality $|V(G)| \geq r + 9$ by induction with respect to r . From Table 1.1 we see that

$$R(r - 2, 3) < r + 8, \quad 5 \leq r \leq 7. \quad (5.2)$$

Obviously, from $G \in H_v(2r; r - 2)$ it follows that $\chi(G) \neq \text{cl}(G)$. Thus, $\alpha(G) \geq 2$. From (5.1) we obtain $|V(G)| \geq r + 8$. This, together with (5.2), implies $|V(G)| > R(r - 2, 3)$ if $5 \leq r \leq 7$. Since $\text{cl}(G) < r - 2$, $\alpha(G) \geq 3$. By the inequality (5.1), $|V(G)| \geq r + 9$, $5 \leq r \leq 7$.

Let $r \geq 8$. Obviously, it suffices to consider only the situation when

$$|V(G)| = F_v(2r; r - 2). \quad (5.3)$$

By (5.3) and Lemma 2.2 we have that

$$G \text{ is a vertex-critical } (r + 1)\text{-chromatic graph}; \quad (5.4)$$

and

$$\text{cl}(G) = r - 3. \quad (5.5)$$

From (5.4) and (5.5) it follows that

$$f(G) = 4. \quad (5.6)$$

We shall consider separately two cases.

Case 1. $|V(G)| < 2r + 1$. By (5.4) and Theorem 2.1 we obtain that

$$G = G_1 + G_2. \quad (5.7)$$

From (5.7), (2.1) and (5.4) obviously it follows that

$$G_i, i = 1, 2 \text{ is a vertex-critical chromatic graph}. \quad (5.8)$$

Let $f(G_1) = 0$. Then, according to (5.8) G_1 is a complete graph. Thus, it follows from (5.7) that $G = K_1 + G'$. It is clear that

$$G \in H_v(2r; r - 2) \Rightarrow G' \in H_v(2r - 1; r - 3).$$

By the induction hypothesis, $|V(G')| \geq r + 8$. Hence, $|V(G)| \geq r + 9$. Let $f(G_i) \neq 0$, $i = 1, 2$. We see from (5.7), (2.3) and (5.6) that $f(G_i) \leq 3$, $i = 1, 2$. By Corollary 4.1 we conclude that

$$|V(G_i)| \geq \chi(G_i) + 2f(G_i), \quad i = 1, 2.$$

Summing these inequalities and using (2.1) and (2.3) we obtain

$$|V(G)| \geq \chi(G) + 2f(G). \quad (5.9)$$

According to (5.4), $\chi(G) = r + 1$. Finally, from (5.9) and (5.6) it follows that $|V(G)| \geq r + 9$.

Case 2. $|V(G)| \geq 2r+1$. Since $r \geq 8$, then $2r+1 \geq r+9$. Hence $|V(G)| \geq r+9$.

Proof of Theorem 1.6(b) By Theorem 1.6(a), $F_v(2r; r-2) \geq r+9$. Therefore, we need to prove the opposite inequality $F_v(2r; r-2) \leq r+9$ if $r \geq 8$. If $r \geq 9$, this inequality follows from Theorem 3.1 ($s = 1$). Let $r = 8$. By $R(6, 3) = 18$ [11] (see also [7]), there is a graph Q such that $|V(Q)| = 17$, $\alpha(Q) = 2$ and $\text{cl}(Q) = 5$. From $|V(Q)| = 17$ and $\alpha(Q) = 2$ obviously it follows that $\chi(Q) \geq 9$. Thus, by (1.3), $Q \xrightarrow{v} (2_8)$. Hence $Q \in H_v(2_8; 6)$ and $F_v(2_8; 6) \leq |V(Q)| = 17$. Theorem 1.6 is proved. \square

Corollary 5.1. *Let G be a graph such that $f(G) \leq 4$. Then*

$$|V(G)| \geq \chi(G) + 2f(G).$$

Proof. If $f(G) \leq 3$, then Corollary 5.1 follows from Corollary 4.1. Let $f(G) = 4$ and $\chi(G) = r+1$, then $\text{cl}(G) = r-3$. Since $\chi(G) \neq \text{cl}(G)$, we have $\text{cl}(G) \geq 2$, and consequently, $r \geq 5$. By (1.3), $G \in H_v(2_r; r-2)$. Using Theorem 1.6(a), we get $|V(G)| \geq r+9 = \chi(G) + 2f(G)$. \square

Let $r \geq 3s+8$. Define

$$\tilde{Q} = K_{r-3s-8} + Q + \underbrace{C_5 + \cdots + C_5}_s,$$

where graph Q is given in the proof of Theorem 1.6(b). Since $\text{cl}(Q) = 5$ and $\chi(Q) \geq 9$, we have by (2.1) and (2.2) that $\text{cl}(\tilde{Q}) = r-s-3$ and $\chi(\tilde{Q}) \geq r+1$. According to (1.3), $\tilde{Q} \in H_v(2_r; r-s-2)$. Thus, $F_v(2_r; r-s-2) \leq |V(\tilde{Q})|$. Since $|V(\tilde{Q})| = r+2s+9$, we obtain the following

Theorem 5.1. *Let r and s be non-negative integers and $r \geq 3s+8$. Then*

$$F_v(2_r; r-s-2) \leq r+2s+9.$$

ACKNOWLEDGEMENTS. This work was supported by the Scientific Research Fund of Sofia University under Contract no. 85/2006.

6. REFERENCES

1. Chvátal, V.: The minimality of the Mycielski graph. *Lecture Notes Math.*, **406**, 1979, 243–246.

2. Dirac, G.: Map colour theorems related to the Heawood colour formula. *J. London Math. Soc.*, **31**, 1956, 460–471.
3. Folkman, J.: Graphs with monochromatic complete subgraphs in every edge coloring. *SIAM J. Appl. Math.* **18**, 1970, 19–24.
4. Gallai, T.: Kritische graphen II, *Publ. Math. Inst. Hungar. Acad. Sci.*, **8**, 1963, 373–395.
5. Gallai, T.: Critical Graphs. In: Theory of Graphs and Its Applications, Proceedings of the Symposium held in Smolenice in June 1963, Czechoslovak Acad. Sciences, Prague, 1964, pp. 43–45.
6. Greenwood, R., A. Gleason: Combinatorial relation and chromatic graphs. *Canad. J. Math.*, **7**, 1955, 1–7.
7. Grinstead, C., S. Roberts: On the Ramsey numbers $R(3, 8)$ and $R(3, 9)$. *J. Comb. Theory*, **B33**, 1982, 27–51.
8. Guta, P.: On the structure of k -chromatic critical graphs of order $k + p$. *Stud. Cerc. Math.*, **50**, 1998, No 3–4, 169–173.
9. Jensen, T., G. Royle: Small graphs with chromatic number 5: a computer research. *J. Graph Theory*, **19**, 1995, 107–116.
10. Kalbfleish, J.: On an Unknown Ramsey Number. *Michigan Math. J.*, **13**, 1966, 385–392.
11. Kery, G.: On a theorem of Ramsey. *Mat. Lapok*, **15**, 1964, 204–224.
12. Kolev, N., N. Nenov: Folkman number $F_e(3, 4; 8)$ is equal to 16. *Compt. rend. Acad. bulg. Sci.*, **59**, 2006, No 1, 25–30.
13. Kolev, N., N. Nenov: New recurrent Inequality on a Class of Vertex Folkman Numbers. In: Proc. 35th Spring Conf. of Union Bulg. Mathematicians, Borovets, April 5–8, 2006, pp. 164–168.
14. Łuczak, T., A. Ruciński, S. Urbański: On minimal vertex Folkman graphs. *Discrete Math.*, **236**, 2001, 245–262.
15. Mycielski, J.: Sur le coloriage des graphes. *Colloq. Math.*, **3**, 1955, 161–162.
16. Nenov, N.: Lower bound for some constants related to Ramsey graphs. *Annuaire Univ. Sofia Fac. Math. Mech.*, **75**, 1981, 27–38 (in Russian).
17. Nenov, N.: On the Zykov numbers and some its applications to Ramsey theory. *Serdica Bulg. Math. Publ.*, **9**, 1983, 161–167 (in Russian).
18. Nenov, N.: The chromatic number of any 10-vertex graph without 4-cliques is at most 4. *Compt. rend. Acad. bulg. Sci.*, **37**, 1984, 301–304 (in Russian).
19. Nenov, N.: On the small graphs with chromatic number 5 without 4-cliques. *Discrete Math.*, **188**, 1998, 297–298.
20. Nenov, N.: On a class of vertex Folkman graphs. *Annuaire Univ. Sofia Fac. Math. Inform.*, **94**, 2000, 15–25.
21. Nenov, N.: A generalization of a result of Dirac. *Ann. Univ. Sofia Fac. Math. Inform.*, **95**, 2001, 59–69.
22. Nenov, N.: On the 3-coloring vertex Folkman number $F(2, 2, 4)$. *Serdica Math. J.*, **27**, 2001, 131–136.
23. Nenov, N.: On a class of vertex Folkman numbers. *Serdica Math. J.*, **28**, 2002, 219–232.
24. Nenov, N.: On the triangle vertex Folkman numbers. *Discrete mathematics*, **271**, 2003, 327–334.

25. Radziszowski, S.: Small Ramsey numbers. *The Electronic Journal of Combinatorics*, Dynamic Survey, version 11, August 1, 2006.

Received on March 31, 2007
In revised form on September 1, 2007

Faculty of Mathematics and Informatics
Sofia University "St. Kliment Ohridski"
5, J. Bourchier Blvd., BG-1164 Sofia
BULGARIA
e-mail: nenov@fmi.uni-sofia.bg